

**Typical Analysis**

NUTRIENT	PER 100 KCAL ME	AS FED	DRY MATTER
Moisture	-	5.5 %	-
Crude Protein	12.3 g	42.0 %	44.4 %
Animal Protein	10.3 g	35.3 %	37.3 %
Vegetable Protein	2.0 g	6.7 %	7.1 %
Crude Fat	4.0 g	13.5 %	14.3 %
Carbohydrates (NFE)	6.6 g	22.5 %	23.8 %
Dietary Starch	4.0 g	13.5 %	14.3 %
Sugars	0.1 g	0.5 %	0.5 %
Crude Fiber	2.3 g	8.0 %	8.5 %
Total Dietary Fiber	4.0 g	13.5 %	14.3 %
Calcium	0.4 g	1.4 %	1.5 %
Phosphorus	0.3 g	1.1 %	1.2 %
Sodium	0.2 g	0.7 %	0.7 %
Chloride	0.3 g	0.9 %	1.0%
Potassium	0.2 g	0.7 %	0.7 %
Magnesium	0.03 g	0.1 %	0.1 %
Copper	0.5 mg	17.0 mg/kg	18.0 mg/kg
Zinc	4.7 mg	160 mg/kg	169 mg/kg
Taurine	90.9 mg	3100 mg/kg	3280 mg/kg
Methionine	0.4 g	1.3 %	1.4 %
Methionine + Cysteine	0.5 g	1.7 %	1.8 %
Total Omega-6 Fatty Acids	0.82 g	2.80%	2.96 %
Total Omega-3 Fatty Acids	0.23 g	0.80%	0.85 %
Omega-6 to Omega-3 Ratio		3.5	

**Specific Functional Ingredients**

L-Carnitine	16.4 mg	560 mg/kg	593 mg/kg
-------------	---------	-----------	-----------

**Calorie Content (Calculated)**

kcal per kg	3410
kcal per lb	1547
Grams per 8 oz Cup	95
kcal per Cup	324

**Digestion Test Results**

Protein, %	82
Fat, %	91

**Metabolizable Energy**

% ME Protein	43
% ME Fat	34
% ME Carbohydrate	23

**Urinary Parameters\***

Urinary pH	6.0 - 6.5
Struvite RSS	< 2.5
Oxalate RSS	< 12

\*Typical values achieved during clinical study when diet was fed as sole nutrition. Values in this range are known to promote a urinary environment unfavorable for the development of struvite and calcium oxalate uroliths.

