

Brief Summary of Prescribing Information for Swine

Before using TULISSIN® 25 (tulathromycin injection) consult the product insert, a summary of which follows:



Tulissin®25
(tulathromycin injection)
Injectable Solution

Antibiotic

25 mg of tulathromycin/mL
For use in swine

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

INDICATIONS:**Swine**

TULISSIN 25 Injectable Solution is indicated for the treatment of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Bordetella bronchiseptica*, *Haemophilus parasuis*, and *Mycoplasma hyopneumoniae*; and for the control of SRD associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, and *Mycoplasma hyopneumoniae* in groups of pigs where SRD has been diagnosed.

DOSAGE AND ADMINISTRATION**Swine**

Inject intramuscularly as a single dose in the neck at a dosage of 2.5 mg/kg (1 mL/22 lb) Body Weight (BW). Do not inject more than 4 mL per injection site.

Table 1. TULISSIN 25 Swine Dosing Guide (25 mg/mL)

Animal Weight (Pounds)	Dose Volume (mL)
4	0.2
10	0.5
15	0.7
20	0.9
22	1.0
25	1.1
30	1.4
50	2.3
70	3.2
90	4.0

Brief Summary of Prescribing Information for Swine

Before using TULISSIN® 100 (tulathromycin injection) consult the product insert, a summary of which follows:



Tulissin®100
(tulathromycin injection)
Injectable Solution

Antibiotic

100 mg of tulathromycin/mL
For use in swine.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

INDICATIONS:**Swine**

TULISSIN 100 Injectable Solution is indicated for the treatment of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Bordetella bronchiseptica*, *Haemophilus parasuis*, and *Mycoplasma hyopneumoniae*; and for the control of SRD associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, and *Mycoplasma hyopneumoniae* in groups of pigs where SRD has been diagnosed.

DOSAGE AND ADMINISTRATION

Inject intramuscularly as a single dose in the neck at a dosage of 2.5 mg/kg (0.25 mL/22 lb) BW. Do not inject more than 2.5 mL per injection site.

Table 1. TULISSIN 100 Swine Dosing Guide. (refer to Table 2 on product insert)

Animal Weight (Pounds)	Dose Volume (mL)
15	0.2
30	0.3
50	0.6
70	0.8
90	1.0
110	1.3
130	1.5
150	1.7
170	1.9
190	2.2
210	2.4
230	2.6
250	2.8
270	3.1
290	3.3

CONTRAINDICATIONS

The use of TULISSIN 25 Injectable Solution is contraindicated in animals previously found to be hypersensitive to the drug.

WARNINGS

FOR USE IN ANIMALS ONLY.

NOT FOR HUMAN USE.

KEEP OUT OF REACH OF CHILDREN.

NOT FOR USE IN CHICKENS OR TURKEYS.

RESIDUE WARNINGS

Swine intended for human consumption must not be slaughtered within 5 days from the last treatment.

PRECAUTIONS**Swine**

The effects of TULISSIN 25 Injectable Solution on porcine reproductive performance, pregnancy, and lactation have not been determined. Intramuscular injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

ADVERSE REACTIONS**Swine**

In one field study, one out of 40 pigs treated with tulathromycin injection (100 mg/mL) at 2.5 mg/kg BW exhibited mild salivation that resolved in less than four hours.

STORAGE CONDITIONS:

Store at or below 30°C (86°F). Use within 45 days of first puncture and puncture a maximum of 30 times. Consider using automatic injection equipment or a repeater syringe. When using a needle or draw-off spike larger than 16 gauge, discard any remaining product immediately after use.

Manufactured for:

Virbac AH, Inc.

P.O. Box 162059, Fort Worth, TX 76161

Made in France

Approved by FDA under ANADA # 200-668

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Virbac AH, Inc. at 1-800-338-3659 or us.virbac.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

TAKE TIME



OBSERVE LABEL DIRECTIONS

©2021 Virbac Corporation. All Rights Reserved.
TULISSIN is a registered trademark of Virbac S.A.

Rev. 06/21

CONTRAINDICATIONS

The use of TULISSIN 100 Injectable Solution is contraindicated in animals previously found to be hypersensitive to the drug.

WARNINGS

FOR USE IN ANIMALS ONLY.

NOT FOR HUMAN USE.

KEEP OUT OF REACH OF CHILDREN.

NOT FOR USE IN CHICKENS OR TURKEYS.

RESIDUE WARNINGS

Swine intended for human consumption must not be slaughtered within 5 days from the last treatment.

PRECAUTIONS**Swine**

The effects of TULISSIN 100 on porcine reproductive performance, pregnancy, and lactation have not been determined. Intramuscular injection can cause a transient local tissue reaction that may result in trim loss of edible tissue at slaughter.

ADVERSE REACTIONS**Swine**

In one field study, one out of 40 pigs treated with tulathromycin injection at 2.5 mg/kg BW exhibited mild salivation that resolved in less than four hours.

STORAGE CONDITIONS:

Store at or below 30°C (86°F). Use within 45 days of first puncture and puncture a maximum of 20 times. Consider using automatic injection equipment or repeater syringe. When using a needle or draw-off spike larger than 16 gauge, discard any remaining product immediately after use.

Manufactured for:

Virbac AH, Inc.

P.O. Box 162059, Fort Worth, TX 76161

Made in France

Approved by FDA under ANADA # 200-669

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Virbac AH, Inc. at 1-800-338-3659 or us.virbac.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

TAKE TIME



OBSERVE LABEL DIRECTIONS

©2021 Virbac Corporation. All Rights Reserved.
TULISSIN is a registered trademark of Virbac S.A.

Rev. 10/21