

Approved by FDA under NADA # 065-492

BIOMOX®

(amoxicillin tablets)

For use in DOGS only.

DESCRIPTION:

BIOMOX tablets are a broad-spectrum, semisynthetic antibiotic which provides bactericidal activity against a wide range of common gram-positive and gram-negative pathogens. Amoxicillin chemically is D-(-) α -amino-p-hydroxybenzyl penicillin trihydrate.

Inactive Ingredients:

Dibasic Calcium Phosphate Dihydrate, Magnesium Stearate, Microcrystalline Cellulose and Sodium Starch Glycolate.

ACTION:

Amoxicillin has bactericidal activity against susceptible organisms similar to that of ampicillin. It acts by inhibiting the biosynthesis of bacterial cell wall mucopeptides. Most strains of the following gram-positive and gram-negative bacteria have demonstrated susceptibility to amoxicillin, both *in vitro* and *in vivo*: nonpenicillinase-producing staphylococci, alpha- and beta-hemolytic streptococci, *Enterococcus faecalis*, *Escherichia coli* and *Proteus mirabilis*. Amoxicillin does not resist destruction by penicillinase; therefore, it is not effective against penicillinase-producing bacteria, particularly resistant staphylococci. Most strains of *Enterobacter* and *Klebsiella* and all strains of *Pseudomonas* are resistant. Amoxicillin may be given without regard to meals because it is stable in gastric acid. It is rapidly absorbed following oral administration and diffuses readily into most body fluids and tissues. It diffuses poorly into the brain and spinal fluid except when the meninges are inflamed. Most of the amoxicillin is excreted in the urine unchanged.

INDICATIONS:

BIOMOX® (amoxicillin tablets) are indicated for treatment of the following infections in dogs when caused by susceptible strains of organisms:

BACTERIAL DERMATITIS due to *Staphylococcus aureus*, *Streptococcus* spp., *Staphylococcus* spp., and *Escherichia coli*.

SOFT TISSUE INFECTIONS

(abscesses, wounds, lacerations) due to *Staphylococcus aureus*, *Enterococcus faecalis*, *Escherichia coli*, *Proteus mirabilis*, and *Staphylococcus* spp.

With all antibiotic therapy, appropriate *in vitro* cultures and sensitivities should be conducted prior to treatment.

CONTRAINDICATIONS:

Use of amoxicillin is contraindicated in animals with a history of an allergic reaction to penicillin.

ADVERSE REACTIONS:

Amoxicillin is a semisynthetic penicillin and, therefore, has the potential for producing allergic reactions. Epinephrine and/or steroids should be administered if an allergic reaction occurs.

WARNINGS:

For use in dogs only.

PRECAUTIONS:

Until adequate reproductive studies are accomplished, BIOMOX® (amoxicillin tablets) should not be used in pregnant or breeding animals.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DOSAGE AND ADMINISTRATION:

The recommended dosage is 5 mg per pound of body weight administered twice daily for 5 to 7 days or 48 hours after all clinical signs have subsided. If no improvement is noted in 5 days, the diagnosis should be reconsidered and therapy changed.

SUPPLY:

BIOMOX® (amoxicillin tablets) are supplied in 50 mg, 100 mg and 200 mg concentrations in bottles of 500 tablets.

Manufactured for:

Virbac AH, Inc.

P.O. Box 162059

Fort Worth, TX 76161

1-800-338-3659

Printed in USA

Rev. -06 for 50 mg and 100 mg; Rev. -07 for 200 mg
05/23

© 2023 Virbac Corporation. All Rights Reserved.
BIOMOX is a registered trademark of the Virbac Group of Companies.